



RECEIVED PCT  
JC10 REC'D PCT/PTO 26 FEB 2002

FORM PTO-1390(modified) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE  
(REV 10-95)

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING  
A FILING UNDER 35 U.S.C., 371

ATTORNEY'S DOCKET NUMBER  
02/23383

U.S. APPLICATION NO. (IF KNOWN), SEE 37 CFR

10/069455

INTERNATIONAL APPLICATION NO.  
PCT/IL00/00510

INTERNATIONAL FILING DATE  
27 AUGUST 2000

PRIORITY DATE CLAIMED  
1 SEPTEMBER 1999

TITLE OF INVENTION

TRICYCLIC COMPOUNDS AND THEIR USES AS ANTIARRHYTHMIC  
ANTIFIBRILLATORY AND DEFIBRILLATORY AGENTS

APPLICANT(S) FOR DO/EO/US


1) Mordechai EREZ 2) Ofra LEVY 3) Ehud KEINAN

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19<sup>th</sup> month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(3)(2)
  - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(3)(2).
7. ☒ A copy of the International Search Report (PCT/ISA/210).
8. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)
  - a. ☒ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☐ have not been made and will not be made.
9. ☐ A translation of the amendments to the claims under PCT Article 19(35 U.S.C. 371(c)(3)).
10. ☒ An unsigned oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 13 to 18 below concern document(s) of information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☐ A FIRST preliminary amendment.  
A SECOND or SUBSEQUENT preliminary amendment.
16. ☐ A substitute specification.
17. ☐ A change of power of attorney and/or address letter.
18. ☐ Certificate of Mailing by Express Mail
19. ☐ Sequence Listing Statement: The sequence listing information recorded in computer readable form is identical to the written (on paper or compact disc) sequence listing.
20. ☐ Other items or information:

U.S. APPLICATION NO. (IF KNOWN) <b>10/069455</b>		INTERNATIONAL APPLICATION NO. <b>PCT/IL00/00510</b>		ATTORNEY'S DOCKET NUMBER <b>02/23383</b>	
20. The following fees are submitted:  <b>BASIC NATIONAL FEE (37 CFR 1.492(a) (1) – (5)):</b>  <input type="checkbox"/> Search Report has been prepared by the EPO or JPO \$ 890 <input checked="" type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1,482) \$ 710 <input type="checkbox"/> No international preliminary examination fee paid to USPTO (37 CFR 1,482) but international search fee paid to USPTO (37 CFR 1,445(a)(2)) \$ 740 <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$ 1040 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) \$ 100  <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>				<b>CALCULATIONS</b> PTO USE ONLY  \$710.00 \$0.00 \$0.00 \$360.00 \$252.00 \$0.00 \$1,322.00 \$661.00 \$661.00 \$0.00 \$661.00 \$0.00 \$661.00 Amount to be refunded: \$ charged \$	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e))					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	40 - 20 =	20	x \$ 18	\$360.00	
Independent claims	6 - 3 =	3	x \$ 84	\$252.00	
Multiple Dependent Claims (check if applicable)			<input type="checkbox"/>	\$0.00	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$1,322.00	
Reduction of 1/2 for filing by small entity, if applicable.			<input checked="" type="checkbox"/>	\$661.00	
<b>SUBTOTAL =</b>				\$661.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f))				+ \$0.00	
<b>TOTAL NATIONAL FEE =</b>				\$661.00	
Fee for recording the enclosed assignment (37 CFR 1.2(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable)			<input type="checkbox"/>	\$0.00	
<b>TOTAL FEES ENCLOSED =</b>				\$661.00	
				Amount to be refunded: \$	
				charged \$	
<input type="checkbox"/> A check in the amount of \$ _____ to cover the above fees is enclosed.  <input checked="" type="checkbox"/> Please charge my Deposit Account No. <b>50-1407</b> in the amount of \$ <b>661.00</b> to cover the above fees. A duplicate copy of this sheet is enclosed.  <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. <b>50-1407</b> . A duplicate copy of this sheet is enclosed.  <b>NOTE: Where an appropriate time limit under 37CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.</b>  SEND ALL CORRESPONDENCE TO:					
SOL SHEINBEIN G.E. EHRLICH (1995) LTD. C/O ANTHONY CASTORINA SUITE 207 2001 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202, USA			<div style="text-align: center;">             _____            SIGNATURE         </div> <div style="text-align: center;">           SOL SHEINBEIN            _____            NAME         </div> <div style="text-align: center;"> <b>25,457</b>            _____            REGISTRATION NUMBER         </div> <div style="text-align: center;"> <b>Feb 24, 2002</b>            _____            DATE         </div>		

1  
TRICYCLIC COMPOUNDS AND THEIR USES AS ANTIARRHYTHMIC  
ANTIFIBRILLATORY AND DEFIBRILLATORY AGENTS

Rec'd PCT/PTO 26 FEB 2002 <sup>fb</sup>FIELD AND BACKGROUND OF THE INVENTION

5           The present invention relates to tricyclic compounds of the 11-oxo-dibenzodiazepin and dibenzoazepin families, and their new therapeutic uses as defibrillating, and/or anti-fibrillatory, and/or anti-arrhythmic and/or anti-ischemic drugs. More particularly, the present invention relates to 5(N-acyl)-derivatives and 5(N- $\beta$ -aminoalcohol)-derivatives of  
10 11-Oxo-10,11-dihydro-dibenzo[b,e] [1,4]diazepin and 10,11-dihydro-dibenzo[b,f]azepin, compositions including same, methods of their synthesis, purification and formulation and their use in prevention and treatment of cardiac disorders, such as, but not limited to, arrhythmia and ventricular fibrillation.

15           Sudden cardiac death is a leading cause of mortality, and ventricular fibrillation (VF) is thought to play a major role in sudden cardiac death [1]. Ventricular fibrillation can be divided into two categories: sustained VF (SVF), which is fatal, unless external defibrillating intervention is practiced, or transient VF (TVF), which terminates spontaneously. Currently, only one  
20 effective approach has been found to terminate SVF once SVF is initiated, which is the application of electrical defibrillation. Electrical defibrillation can be applied either externally or internally, by implantation. However, this approach has a number of disadvantages. For example, electrical defibrillation must be applied immediately to be effective, yet may not be sufficient, and  
25 may even cause damage. In addition, an implanted defibrillator requires invasive treatment. Thus, artificial defibrillation is not a cure, and does not prevent reoccurrence of VF.

          Antiarrhythmic drugs constitute an alternative, preferable approach, as they are aimed at preventing initiation of VF by decreasing the incidence of  
30 ventricular arrhythmias that can lead to VF [2]. In addition, certain drugs, such as bretylium, have been shown to transform SVF into TVF [3]. The

effectiveness of this treatment, however, is limited, since various mechanisms can be involved in the initiation of VF, such that antiarrhythmic drugs are unlikely to absolutely eliminate arrhythmias and totally prevent VF initiation. Furthermore, recent surveys (such as the Cardiac Arrhythmia Suppression Trial, CAST, II and I [4,5]) have clearly shown limitations to this approach. Thus, a new cardiac protective therapy is needed. For this reason, a new approach has been proposed, to use a new class of antiarrhythmic drugs [6], which can enhance spontaneous termination of VF, once it occurs. In several animal species, and even, though rarely, in humans, VF can revert spontaneously into sinus rhythm, resulting in the non-fatal TVF. It has been previously found that several factors contribute to the ability to self-defibrillate. For example, self-defibrillation is a normal feature of young mammals, but this ability decreases with age [7]. Such spontaneous defibrillation requires a relatively high degree of intercellular synchronization [8], and is enhanced by increased sympathetic activity. Thus, treatments with compounds that elevate extraneuronal catecholamine levels in the heart enhance self-defibrillation and administration of  $\beta$ -adrenergic blockers abolishes this activity [9].

In order to design and synthesize new, more potent and selective defibrillatory drugs, it has been found that certain dibenzoazepins (imipramine, desipramine, maprotiline and bonnecore) and phenothiazins (chlopromazin, moricizine and trifluoperazin), induce self-defibrillation and increase the threshold for electrical fibrillation [10, 11]. Moreover, tricyclic antidepressants, in addition to their antiarrhythmic and defibrillating effects, have the ability to decrease the ischemic area in the heart following coronary occlusion [12]. However, these cardio-protective effects of the compounds were expressed when relatively high doses were used, resulting in a low therapeutic index.


There is thus a need for, and it would be useful to have, pharmaceutically effective compounds which are safe and useful for the

treatment of ventricular fibrillation, particularly for the induction of TVF once SVF has been initiated, and for both treating and preventing pathological conditions associated with VF.

## 5 SUMMARY OF THE INVENTION

The present invention relates to 5(N-acyl)-derivatives and 5(N-β-aminoalcohol)-derivatives of 11-Oxo-10,11-dihydro-dibenzo [b,e][1,4] diazepin and 10,11-dihydro-dibenzo[b,f]azepin, compositions including same, methods of their synthesis, purification and formulation and their use in  
10 prevention and treatment of cardiac disorders. It is shown herein for the first time that these new tricyclic compounds and some previously known tricyclic compounds have been synthesized and have been shown to have substantial activity as chemical defibrillating agents.

According to the present invention, there is provided a compound  
15 having a general formula (I):



wherein A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched  
20 alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>C\*HOHCH<sub>2</sub>NR'R'' (the chiral C, which is marked by \* can be the R anantiomer, the S anatiomer, be in a racemic mixture or in any other ratio between the R and S anantiomers), wherein *n* is  
25 an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl, and sulfonylamide; *q* and *t* are each an integer independently selected from 1-4; and pharmaceutically acceptable salts

thereof.

Preferably,  $R_1$  is selected from the group consisting of  $\beta$ -amino-alcohol,  $C(=O)(CH_2)_nNR'R''$  and  $(CH_2)_nCHOHCH_2NR'R''$  wherein  $n$  is an integer and further wherein the chiral carbon atom can be the R or S anantiomer, a racemic mixture thereof or a mixture of any ratio thereof; and  $R'$ ,  $R''$  are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl; and pharmaceutically acceptable salts thereof. Examples of suitable alkyl moieties include but are not limited to *iso*-propyl, *iso*-butyl, *tert*butyl and *sec*-butyl. Most preferably,  $R''$  is *iso*-propyl.

According to a preferred embodiment of the present invention, A and B are each an alkyl chain, more preferably a CH moiety,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are each hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$  and  $(CH_2)_nCHOHCH_2NR'R''$ ,  $n$  being 0-5. Preferably, R and  $R'$  are each hydrogen and  $R''$  is selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl and *tert*-butyl.

According to a preferred embodiment of the present invention A is  $CR_2R_3$  or  $C=O$  and B is  $CR_4R_5$ ; R,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$  and  $(CH_2)_nCHOHCH_2NR'R''$ ,  $n$  being 0-5.

## DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention relates to a group of compounds, based upon the general backbone structure of tricyclic antidepressants, namely, 11-oxo-dibenzodiazepins and dibenzoazepins N-substituted at the 5 position, as well as to pharmaceutical compositions of these compounds and to their use in the treatment and prevention of ventricular fibrillation and ischemic damage by local or systemic application. More specifically, these compounds are demonstrated to have a defibrillating effect on ventricular fibrillation, once it actually occurs.


Previous studies, aimed at characterizing the structure-activity relationships of dibenzoazepins and phenothiazines accountable to improved defibrillating activity, have indicated the relevance of several structural features of the backbone of these molecules for their efficacy against ventricular fibrillation. First, activity of the tricyclic compounds correlated with higher (more bent) dihedral angle between the two benzo rings in the tricyclic skeleton. This angle is also considered to play a detrimental role in the type of central nervous system activity of these compounds, as well as their different selectivity towards catecholamine reuptake systems (such as noradrenaline, serotonin and dopamine). Second, compounds with 5-N-substituted secondary aminoalkyl side chains (such as desipramine) exhibit higher defibrillatory activity, as opposed to tertiary alkylamine (such as in imipramine) and lastly, transition from aminoalkyl to aminoacyl in the side chain at position 10 of the phenothiazine tricyclic nucleus results in an increase in antiarrhythmic activity and decrease in psychotropic activity [10].

The development of the disclosed class of tricyclic compounds was based on the rationalized design of new compounds, aimed at more focused and selective activity as chemical defibrillators, as well as identifying additional molecules which are able to overcome shortcomings of the presently utilized approaches (electrical defibrillation, antiarrhythmic drugs, described above), and which are able to convert the fatal sustained ventricular fibrillation to the non-fatal transient one.

The experiments described below in the Examples section demonstrate that the disclosed compounds are indeed effective in transforming the potentially fatal VF type, SVF, to the spontaneously-defibrillating type, TVF. In addition, the preferred structure is indicated in terms of a structure-activity-relationship, indicating the importance of certain structural elements in obtaining potent, selective therapeutic activity, and minimizing untoward side effects. Thus, a compound according to the present invention includes derivatives of 5-(N-alkyl), 5-(N-acyl) and 5-(N- $\beta$ -aminoalcohol) dibenzoazepins of the

6

general formula (II):




[II]

wherein  $R_1$  is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl,  $C(=O)(CH_2)_nNR'R''$  or  $(CH_2)_nCHOHCH_2NR'R''$ , wherein  $n$  is an integer;  $R_Q$ ,  $R_T$ ,  $R'$ , and  $R''$  are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and  $t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

Preferably,  $R_1$  is selected from the group consisting of  $\beta$ -amino-alcohol,  $C(=O)(CH_2)_nNR'R''$ , and  $(CH_2)_nCHOHCH_2NR'R''$  wherein  $n$  is an integer and  $R'$ ,  $R''$  are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl; and pharmaceutically acceptable salts thereof. Examples of suitable alkyl moieties include but are not limited to *iso*-propyl, *iso*-butyl, *tert*-butyl and *sec*-butyl. Most preferably,  $R''$  is *iso*-propyl. Also most preferably,  $R_Q$  and  $R_T$  are each a hydrogen, and  $q$  and  $t$  are each 4.

The compounds of the present invention are also useful as an active ingredient in a composition for treating or preventing a cardiac disorder, such as ventricular fibrillation, featuring a pharmaceutically effective amount of a tricyclic compound in combination with a pharmaceutically acceptable carrier, in which the tricyclic compound is of a general formula (I):





wherein A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', wherein *n* is an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl, and sulfonylamide; *q* and *t* are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

Preferably, R<sub>1</sub> is selected from the group consisting of β-amino-alcohol, C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', and (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'' wherein *n* is an integer and R', R'' are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl; and pharmaceutically acceptable salts thereof. Examples of suitable alkyl moieties include but are not limited to *iso*-propyl, *iso*-butyl, *tert*-butyl and *sec*-butyl. Most preferably, R'' is *iso*-propyl.

According to a preferred embodiment of the present invention, A and B are each an alkyl chain, more preferably a CH moiety, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are each hydrogen, and R<sub>1</sub> is C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'' and (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', *n* being 0-5. Preferably, R and R' are each hydrogen and R'' is selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl and *tert*-butyl.

According to a preferred embodiment of the present invention A is CR<sub>2</sub>R<sub>3</sub> or C=O and B is CR<sub>4</sub>R<sub>5</sub>, R, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each a hydrogen, and R<sub>1</sub> is C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'' and (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', *n* being 0-5. More preferably, *n* is 1 or 2.

Hereinafter, the term "tricyclic compound" refers to 5-(N-acyl) or 5-(N-alkylβaminoalcohol)-derivatives or 5-(N-alkyl)-derivatives of

10,11-dihydro-dibenzo[b,f]azepin or 5-(N-acyl) or  
5-(N-alkylβaminoalcohol)-derivatives or 5-(N-alkyl)- derivatives of  
11-Oxo-10,11-dihydro-dibenzo[b,e][1,4] diazepin compounds of the present  
invention.

5 Hereinafter, the term "derivative" refers to the result of a chemically  
altering, modifying or changing a molecule or a portion thereof, such that the  
molecule either maintains or increases its functionality.

Hereinafter, the term "pharmaceutically acceptable carrier" refers to a  
carrier or a diluent that does not cause significant irritation to an organism and  
10 does not abrogate the biological activity and properties of the administered  
compound.

Hereinafter, the term "pharmaceutically effective amount" refers to that  
amount of the compound being administered which will relieve to some extent  
one or more of the symptoms of the disorder being treated.

15 The composition of the present invention may be optionally and  
preferably designed for topical application such as a slow release or  
transdermal patch of the tricyclic compound. The composition for slow  
release includes particles including a slow release carrier (typically, a  
polymeric carrier), and the tricyclic compound. Slow release biodegradable  
20 carriers are well known in the art. These are materials that may form particles  
that may capture therein an active compound(s) and slowly degrade/dissolve  
under a suitable environment (e.g., aqueous, acidic, basic, etc.) and thereby  
degrade/dissolve in body fluids and release the active compound(s) therein.

Specifically, a slow release formulation or a transdermal patch of the  
25 tricyclic compound can be used in patients prone to cardiac disorders, such as  
arrhythmias, or with a known history of arrhythmic or fibrillatory episodes.

One particularly preferred embodiment of the present invention is  
parenteral administration of the tricyclic compound, for example  
intravenously. Further according to the present invention there is provided a  
30 method for treating or preventing a cardiac disorder, such as ventricular

fibrillation, in a subject, by administering a pharmaceutically effective amount of a tricyclic compound of a general formula (I):



wherein A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', wherein *n* is an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl, and sulfonylamide; *q* and *t* are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

Preferably, R<sub>1</sub> is selected from the group consisting of β-amino-alcohol, C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', and (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'' wherein *n* is an integer and R', R'' are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl; and pharmaceutically acceptable salts thereof. Examples of suitable alkyl moieties include but are not limited to *iso*-propyl, *iso*-butyl, *tert*butyl and *sec*-butyl. Most preferably, R'' is *iso*-propyl.

According to a preferred embodiment of the present invention, A and B are each an alkyl chain, more preferably a CH moiety, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are each hydrogen, and R<sub>1</sub> is C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'' and (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', *n* being 0-5. Preferably, *n* is 1 or 2. Preferably, R and R' are each hydrogen and R'' is selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl and *tert*-butyl.

According to a preferred embodiment of the present invention A is  $CR_2R_3$  or  $C=O$  and B is  $CR_4R_5$ ;  $R$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$  and  $(CH_2)_nCHOHCH_2NR'R''$ ,  $n$  being 0-5.

Preferably, this method of treatment with the compositions of the present invention is used as an adjunct or additive therapy for patients with implanted defibrillators, in order to combine the electrical defibrillation with the chemical defibrillation. The utility of such a combination has been previously shown [M.T. Neuman, *PhD Thesis*, 1995, Tel Aviv University, Tel Aviv, Israel].

Herein, the term "treating" includes substantially inhibiting or completely stopping episodes of cardiac dysfunction, such as ventricular fibrillation, in a subject.

Herein, the term "preventing" refers to a method for barring a subject from exhibiting symptoms of a cardiac disorder, such as ventricular fibrillation, or alternatively for at least reducing the likelihood and/or severity of such symptoms arising in the subject.

Further according to the present invention there is provided a method of locally treating a disorder of a tissue of a subject comprising the step of locally applying the herein above described composition to the tissue. The method includes the steps of applying the composition to an implant and inserting the implant into a tissue, such as cardiac tissue. Alternatively, the composition can be applied to a transdermal patch, which is applied to the skin for system absorption.

Also according to the present invention, there is provided a method for preventing ischemia in the cardiac tissue of a subject, by administering the above-referenced compound of the present invention to the subject.

A precise understanding of the mechanism by which the tricyclic compounds of the present invention cause such chemical defibrillation is not required in order to practice the present invention. However, while not wishing to be bound to any particular mechanism or theory, it is believed that increased catecholamine levels, induced by the compounds of the present invention, are

involved in the process. Several lines of evidence support this hypothesis. For example, the evaluation of intra- and inter-species differences for the ability to defibrillate spontaneously have indicated the central role of cardiac autoregulation. In particular, TVF appears in animals with predominantly sympathetic autoregulation, while SVF appears in animals with predominantly vagal autoregulation. Within members of the same species, the ability to defibrillate spontaneously is a normal feature of young mammals, and this ability decreases with age. Respectively, cardiac autoregulation in young mammals is dominantly sympathetic and turns to a vagal predominance with age.

Furthermore, administration of either  $\beta$ -adrenergic blockers (e.g., propranolol or pindolol) or a parasympathomimetic agonist (e.g., acetylcholine or metacholine) in mammals exhibiting TVF, prolongs the duration of TVF and even transforms it into SVF. Lastly, self-defibrillation requires a relatively high degree of intercellular synchronization, which may be enhanced by elevated catecholamine levels. Thus, compounds which are known to elevate extraneuronal catecholamine levels in the heart, such as dibenzoazepins and phenothiazines, were identified to enhance ventricular self-defibrillation, an effect abolished by co-administration of  $\beta$ -adrenergic blockers [9, 14].

Furthermore, in a recent publication, the efficacy in defibrillating activity of dibenzoazepins and phenothiazines was directly related to their ability to inhibit noradrenaline uptake [10]. Alternatively, the compound, 11-Oxo-10,11-dihydro-5-(N-methyl)-propylaminodibenzo[b,e] [1,4] diazepin, was previously suggested for application as a muscarinic receptor antagonist in PCT Application No. WO 91/10654], supporting the role of sympathetic predominance in potential defibrillating activity.

Thus, further according to the present invention there is provided a method of treating or preventing a cardiac disorder in a subject, by inducing cardiac sympathetic activity by administering the tricyclic compounds of the present invention as hereinabove described.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

### EXAMPLES

Reference is now made to the following examples, which together with the above descriptions illustrate the invention in a non-limiting fashion. The following protocols and experimental details are referenced in the Examples that follow.


#### Example 1

##### Synthesis of 10,11-dihydro-5-[(3-N-alkyl)1-oxopropyl]

##### 5-H-dibenzo[b,f]azepin monohydrochloride compounds

**Synthesis of 10,11-dihydro-5-[(3-chloro)1-oxopropyl]-5H-dibenzo[b,f]azepin:** The synthesis was performed as known in the literature (Schindler W.; Hafliger F.; *Helvetica Chima Acta* 1954, **59**, 472-483). To a solution of 5H-10,11-dihydro-dibenzo[b,f] azepin (0.85 gr., 4.359 mmol) in benzene (50 ml), 0.5 ml 3-chloropropionyl chloride (5.225 mmol) was added dropwise. The mixture was refluxed for 3 h, the solvent was removed under reduced pressure and the residue was washed with 5 % aq. HCl, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub> (20 ml). The resulting organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, and the crude product was purified by flash chromatography (silica gel, hexane-ethyl acetate 8:2), resulting in 1.133 gr. (91 % yield) of 10,11-dihydro-5-[(3-chloro)1-oxopropyl]-5-H-dibenzo[b,f]azepin, in the form of white crystals. NMR (CDCl<sub>3</sub>): 7.16 (bs, 8H), 3.82 (m, 2H), 3.46 (m, 1H), 2.85 (m, 2H), 2.51 (m, 1H). MS (CI): 286 (M<sup>+</sup>).

Scheme 1 below illustrates the synthesis and structure of 10,11 dihydro-5-[(3-chloro)1-oxopropyl] 5-H-dibenzo[b,f]azepin:




*Synthesis of 10,11-dihydro-5-[(3-N-alkyl)1-oxopropyl]5-H-dibenzo[b,f]azepin monohydrochloride:*

Synthesis of 10,11-dihydro-5-[(3-N-alkyl)1-oxopropyl]5-H-dibenzo[b,f]azepin derivatives was achieved by reaction of the 10,11-dihydro-5-[(3-chloro)1-oxopropyl]5-H-dibenzo[b,f]azepin as a starting material with the corresponding alkyl amine. The succeeding derivatives, wherein the 3-N- substituted alkyl is a methyl, an ethyl or an iso-propyl, (compounds 15a-c) were prepared following the below-described general procedure: A well-stirred suspension of 10,11-dihydro-5-[(3-chloro)-1-oxopropyl]-5-H-dibenzo [b,f]azepin (1.71 gr., 6.0 mmol) in ethanol (50 ml) was warmed to 65 °C. Alkylamine (10 mmol) was added dropwise. The mixture was stirred for 1 h at 65 °C, and then allowed to cool to room temperature. The mixture was washed with 5 % aq. potassium bicarbonate and extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 ml). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and the solvent was removed under reduced pressure. The crude product was recrystallized from EtOAc at 0 °C. The precipitate was filtered to result in the appropriate free base. The solid was then dissolved in toluene and HCl was bubbled till a precipitate formed. The precipitate was filtered and dried under reduced pressure resulting in the appropriate monohydrochloride salt. Alternatively, wherein bulkier alkyl groups were used for substituting the 3-N alkyl, such as *tert*-butyl, sec-butyl, iso-butyl and benzyl (compounds 15d-g), the following procedure was used: A solution of 10,11-dihydro-5-[(3-chloro)1-oxopropyl]-5-H-dibenzo[b,f] azepin (0.285 gr., 1.0 mmol) in *iso*-propanol (15 ml) and monoamine (15 ml) was refluxed overnight. The solvent was removed under reduced pressure, and the

crude product was recrystallized from EtOAc at 0 °C. The solid was dissolved in toluene and HCl was bubbled till a precipitate was formed. The precipitate was filtered and dried under reduced pressure resulting in the appropriate monohydrochloride salt.

Scheme 2 below illustrates the synthesis and general structure of 10,11-dihydro-5-[(3-N-alkyl)1-oxopropyl]-5-H-dibenzo[b,f]azepin monohydrochloride of both synthetic routes:




SCHEME 2 (15)

10

(i) 10,11-dihydro-5-[(3-N-methyl)-1-oxopropyl]-5-H-dibenzo[b,f]azepin monohydrochloride: The synthesis was performed according to the above synthetic scheme. 1.126 gr. (67 % yield) of light yellow solid was obtained. <sup>1</sup>H NMR (D<sub>2</sub>O): δ 6.87 (brm, 8H), 3.15 (s, 1H), 2.82 (brm, 4H), 2.47 (s, 3H), 2.24 (brm, 3H). FAB/MS 281 (M<sup>+</sup>).

15

Scheme 3 below illustrates the structure of 10,11-dihydro-5-[(3-N-methyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:



20

SCHEME 3 (15 a)


25

(ii) 10,11-dihydro-5-[(3-N-ethyl)1-oxopropyl]5-H-dibenzo[b,f]azepin monohydrochloride: The synthesis was performed according to the above synthetic scheme. 1.129 gr. (64 % yield) of light yellow solid was obtained. <sup>1</sup>H NMR (D<sub>2</sub>O): δ 6.89 (brm, 8H), 3.12 (s, 1H), 2.84 (brm, 7H), 2.27



(brm, 2H), 1.02 (t, J=7.4, 3H). FAB/MS 295 ( $M^+$ ).


Scheme 4 below illustrates the structure of 10,11-dihydro-5-[(3-N-ethyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:



SCHEME 4 (15 b)

(iii) 10,11-dihydro-5-[(3-N-iso-propyl)1-oxopropyl]5-H-dibenzo  
[b,f]azepin monohydrochloride: The synthesis was performed according to  
 the above synthetic scheme. 1.219 gr. (66 % yield) of light yellow solid was  
 obtained, with mp 199.1 °C.  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ ):  $\delta$  6.97 (brm, 8H), 3.14 (m, 1H),  
 2.97 (m, 4H), 2.67 (m, 1H), 2.42 (m, 1H), 2.38 (m, 2H), 1.08 (d, J=6.6, 6H).  
 FAB/MS 309 ( $M^+$ ).

Scheme 5 below illustrates the structure of 10,11-dihydro-5-[(3-N-*iso*-propyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:




SCHEME 5 (15 c)

(iv) 10,11-dihydro-5-[(3-N-tert-butyl)1-oxopropyl]5-H-dibenzo  
[b,f]azepin monohydrochloride: The synthesis was performed according to  
 the above synthetic scheme. 280 mgr. (87 % yield) was obtained.  $^1\text{H}$  NMR  
 ( $\text{CDCl}_3$ ):  $\delta$  7.24 (m, 8H), 3.39 (m, 2H), 2.85 (m, 5H), 2.51 (m, 1H), 1.26 (m,  
 9H). FAB/MS 323 ( $M^+$ ).

Scheme 6 below illustrates the structure of 10,11-dihydro-5- [(3-N-*tert*-butyl)1-oxopropyl]5-H-dibenzo[b,f]azepin monohydrochloride:

16




SCHEME 6 (15 d)

(v) 10,11-dihydro-5-[(3-N-sec-butyl)1-oxopropyl]5-H-dibenzo

[b,f]azepin monohydrochloride: The synthesis was performed according to the above synthetic scheme. 241 mg. (75 % yield) was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.25 (m, 8H), 3.37 (m, 2H), 2.97 (m, 5H), 2.79 (m, 2H), 1.52 (m, 2H), 0.99 (m, 6H). FAB/MS 323 (M<sup>+</sup>).

Scheme 7 below illustrates the structure of 10,11-dihydro-5-[(3-N-sec-butyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:




SCHEME 7 (15 e)

(vi) 10,11-dihydro-5-[(3-N-iso-butyl)1-oxopropyl]5-H-dibenzo

[b,f]azepin monohydrochloride: The synthesis was performed according to the above synthetic scheme. 264 mg. (82 % yield) was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.23 (m, 8H), 3.35 (m, 2H), 2.82 (m, 4H), 2.47 (m, 2H), 1.84 (m, 1H), 0.95 (d, J=6.6 Hz, 3H), 0.89 (d, J=6.6 Hz, 3H). FAB/MS 323 (M<sup>+</sup>).


Scheme 8 below illustrates the structure of 10,11-dihydro-5- [(3-N-iso-butyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:

17


**SCHEME 8 (15 f)****(vii) 10,11-dihydro-5-[(3-N-benzyl)1-oxopropyl]5-H-dibenzo**

**[b,f]azepin monohydrochloride:** The synthesis was performed according to the above synthetic scheme. 367 mg. (67 % yield) was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.23 (m, 8H), 3.35 (m, 2H), 2.82 (m, 4H), 2.47 (m, 2H), 1.84 (m, 1H), 0.95 (d, J=6.6 Hz., 3H), 0.89 (d, J=6.6 Hz, 3H). FAB/MS 357 (M<sup>+</sup>).

Scheme 9 below illustrates the structure of 10,11-dihydro-5- [(3-N-benzyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin monohydrochloride:


**SCHEME 9 (15 g)**

Scheme 9a describes the synthesis of the β-amino alcohol derivative according to the present invention, which was performed in two stages.

**SCHEME 9a**

**10,11-dihydro-5-[N-2 methyloxirane]5-H-dibenzo[b,f]azepin (20).****20r, 20s (Stage I):**

To a solution of 5H-10,11-dihydro-dibenzo[b,f] azepin (1.320 g, 6.711 mmol, see Scheme 1, above) in benzene (50 ml) was added NaNH<sub>2</sub> (0.5 gr). The mixture was refluxed for 2 hr and epichlorohydrin (1 ml, 12.810 mmol) was added dropwise. The mixture was refluxed for 6 hr. The solvent was removed under reduced pressure, and the residue was washed with 5% aq HCl, and extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 ml). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and the crude product was purified by flash chromatography (silica gel, hexane-ethyl acetate 9.5:0.5) to give **20**, (0.75 g, 45%) in the form of white powder. <sup>1</sup>H NMR (chloroform): 7.07 (bs, 4H), 6.93 (m, 2H), 6.73 (bt, 2H) 3.91 (m, 2H), 3.16 (m, 4H), 3.05 (m, 1H), 2.67 (d, 1H), 2.55 (d, 1H). MS (CI): 251 (M<sup>+</sup>). (Scheme 9b)

**SCHEME 9b**

**10,11-dihydro-5-[N-β iso-propylamino-2propanol)]5-H-dibenzo[b,f]azepin monohydrochloride, 25 bi, 25 bis, 25 bir (Stage II)**

**General procedure:** A suspension of **20** (1.5 g, 5.0 mmol) in *iso*-propanol (50 ml) was warmed to 30 °C, *iso*-propyl amine (10 mmol) was added dropwise, the mixture was stirred overnight at this temperature and then allowed to cool to room temperature. The solvent was removed under reduced pressure, and the crude product was recrystallized from ethyl acetate at 0 °C. The precipitation was filtered to give the appropriate free base. The solid was dissolved in toluene and HCl was bubbled until a precipitate reappeared. The precipitation was filtered and dried under reduced pressure to give 25 bi salt.

<sup>1</sup>H NMR (chloroform): 7.07 (bs, 4H), 6.93 (m, 2H), 6.73 (bt, 2H) 3.79 (m, 1H), 3.15 (s, 4H), 2.78 (m, 1H), 2.72 (d, 2H), 2.55 (d, 2H), 1.01 (d, 6H). MS (CI): 311 (M<sup>+</sup>).

### Example 2


5

#### Synthesis of 11-Oxo-10,11-dihydro-5-

#### [(N-alkyl)1oxopropyl]-5-H-dibenzo[b,e][1,4]diazepin monohydrochloride compounds

*Synthesis of 11 - Oxo - 10, 11 - dihydro - 5H - dibenzo [b,e] [1,4] diazepin:* The synthesis was performed according to the below synthetic  
 10 scheme. A suspension of 2-chlorobenzoic acid (5.08 gr., 31.81 mmol), o-phenylenediamine (3.46 gr., 31.48 mmol), copper powder (2.14 gr., 31.7 mmol) and molecular sieves (3A) in chlorobenzene (100 ml) was vigorously stirred at 130 °C for 8 h. The hot mixture was rapidly filtered and the filtrate was concentrated under reduced pressure. The solid precipitate was collected  
 15 by filtration and then recrystallized from EtOH resulting in 11-Oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin 5.48 gr. (82 % yield) in the form of bright yellow powder. <sup>1</sup>H NMR (acetone): 8.89 (s, 1H), 7.80 (d, J=7.7 Hz., 1H), 7.33 (t, J=7.3 Hz., 1H), 7.07 (m, 6H). FAB/MS: 211 (M<sup>+</sup>).

Scheme 10 below illustrates the synthesis and structure of  
 20 11-Oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin monohydrochloride:



25


SCHEME 10 (1)

#### Synthesis of 11-Oxo-10,11-dihydro-5-[(3-chloro)1-oxopropyl]

5-H-dibenzo[b,e][1,4]diazepin: The synthesis was performed according to the below synthetic scheme. To a solution of 11-Oxo-  
 30 10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin (0.376 gr., 1.79 mmol) in dry

THF, (10 ml) NaNH<sub>2</sub> (0.132 gr., 3.39 mmol) was added. The mixture was cooled to 0 °C, and 0.2 ml 3-chloropropionyl chloride (2.09 mmol) was added dropwise. The mixture was stirred for 30 min at 0 °C, allowed warming to room temperature, washed with 5 % aq. potassium bicarbonate and extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 ml). The resulting organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and the solvent was removed under reduced pressure. The crude product was crystallized from ether, resulting in 0.467 gr. (87 % yield) of 11-Oxo-10,11-dihydro-5-[(3-chloro)1-oxopropyl] 5-H-dibenzo[b,e] [1,4]diazepin in the form of a yellow solid, mp 241-242 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.78 (s, 1H), 7.98 (s, 1H), 7.59 (s, 1H), 7.4-7.26 (m, 6H), 3.77 (m, 4H). FAB/MS: 301 (M<sup>+</sup>).

Scheme 11 below illustrates the synthesis and structure of 11-Oxo-10,11-dihydro-5-[(3-chloro)1-oxopropyl] 5-H-dibenzo[b,e] [1,4] diazepin :




SCHEME 11 (6)

**Synthesis of 11-Oxo-10,11-dihydro-5-[(N-alkyl)1-oxopropyl]5-H-dibenzo[b,e] [1,4]diazepin monohydrochloride:** The synthesis of the below described 11-Oxo-10,11-dihydro-5-[(N-alkyl)1-oxopropyl]5-H-dibenzo[b,e][1,4]diazepin hydrochloride compounds, 11a-c, was performed by reacting 11-Oxo-10,11-dihydro-5-[(3-chloro)1-oxopropyl]-5-H-dibenzo[b,e][1,4]diazepin, as a starting material, with the appropriate alkyl amine. A well-stirred suspension of 11-Oxo-10,11-dihydro-5-[(3-chloro)1-oxopropyl]-5-H-dibenzo[b,e][1,4]diazepin (1.5 gr., 5.0 mmol) in ethanol (50 ml) was warmed to 65 °C. Alkylamine (10 mmol) was added dropwise. The mixture was stirred for 1 h at 65 °C, and then allowed to cool to room temperature. The mixture was then washed with 5 % aq. potassium

bicarbonate and extracted thrice with  $\text{CH}_2\text{Cl}_2$  (20 ml). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and the solvent was removed under reduced pressure. The crude product was recrystallized from EtOAc at  $0^\circ\text{C}$ . The precipitate was filtered to result in the appropriate free base. The solid was then dissolved in toluene and HCl was bubbled till a precipitate formed. The precipitate was filtered and dried under reduced pressure resulting in the appropriate hydrochloride salt.


Scheme 12 below illustrates the synthesis and general structure of 11-Oxo-10,11-dihydro-5-[(N-alkyl)1-oxopropyl] 5-H-dibenzo[b,e] [1,4]diazepin monohydrochloride:



SCHEME 12 (11)


(i) *Synthesis of 11-Oxo-10,11-dihydro-5-[(N-methyl)1-oxopropyl] 5-H-dibenzo[b,e] [1,4]diazepin monohydrochloride:* The synthesis was performed according to the above synthetic scheme. The compound, in the form of a light-pink solid was obtained, 627 mg. (44 % yield), with mp=223-225  $^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ ):  $\delta$  7.33 (br m, 8H), 3.06 (t,  $J=6$  Hz, 2H), 2.9 (m, 1H), 2.48 (s, 3H), 2.36 (br m, 1H). FAB/MS 297 ( $\text{M}^+$ ).

Scheme 13 below illustrates the structure of 11-Oxo-10,11-dihydro-5-[(3-N-methyl)1-oxopropyl] 5-H-dibenzo[b,e][1,4]diazepin monohydrochloride:



SCHEME 13 (11 a)

(ii) *Synthesis of 11-Oxo-10,11-dihydro-5-[(N-ethyl)1-oxopropyl]-5-H-dibenzo[b,e][1,4]diazepin monohydrochloride:* The synthesis was performed according to the above synthetic scheme. The compound, in the form of light-yellow solid, was obtained, 1.023 gr. (66 % yield), with mp 243-245 °C. <sup>1</sup>H NMR (D<sub>2</sub>O): 7.40 (br m, 8H), 3.08 (t, J=6.1 Hz., 2H), 2.86 (m, 3H), 2.38 (m, 1H), 1.06 (t, J=7.3, 3H). FAB/MS: 311 (m<sup>+</sup>). Scheme 14 below illustrates the structure of 11-Oxo-10,11-dihydro - 5 - [(N-ethyl)1-oxopropyl] - 5 - H-dibenzo [b,e] [1,4] diazepin hydrochloride:



15

SCHEME 14 (11 b)

(iii) *Synthesis of 11-Oxo-10,11-dihydro-5-[(N-iso-propyl)-1-oxopropyl]-5-H-dibenzo[b,e] [1,4]diazepin monohydrochloride:* The synthesis was performed according to the above synthetic scheme. The compound, in the form of light-orange solid, was obtained, 791 mgr. (49 % yield). <sup>1</sup>H NMR (D<sub>2</sub>O): 7.35 (br m, 8H), 3.05 (br m., 3H), 2.85 (m, 1H), 2.33


20



23

(m, 1H), 1.08 (t, J=7.3, 3H). FAB/MS: 323 (m<sup>+</sup>).

Scheme 14 below illustrates the structure of 11-Oxo-10,11-dihydro-5-[(N-*iso*-propyl) 1-oxopropyl] 5-H-dibenzo [b,e] [1,4] diazepin monohydrochloride:



SCHEME 15(11 c)

### Example 3


Synthesis of 10,11-dihydro-5-[N-β isopropylamino- 2propanol] 5-H-dibenzo [b,f]azepin (25bi). 25bir, 25bis monohydrochloride compounds

#### 10,11-dihydro-5-[N-2 methyloxirane]5-H-dibenzo[b,f]azepin (20).

**20r, 20s:** To a solution of **5** (1.320 g, 6.711 mmol) in benzene (50 mL) was added NaNH<sub>2</sub> (0.5 gr). The mixture was refluxed for 2 hr and the appropriate epichlorohydrin (1 mL, 12.810 mmol) was added dropwise (racemic to obtain **20**, R enantiomer to obtain **20r** and S enantiomer to obtain **20s**. The mixture was refluxed for 6 hrs. The solvent was removed under reduced pressure, and the residue was washed with 5% aq HCl, and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 20 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and the crude product was purified by flash chromatography (silica gel, hexane-ethyl acetate 9.5:0.5) to give **20**, (0.75 g, 45%) in the form of white powder. <sup>1</sup>H NMR (chloroform): 7.07 (bs, 4H), 6.93 (m, 2H), 6.73 (bt, 2H) 3.91 (m, 2H), 3.16 (m, 4H), 3.05 (m, 1H), 2.67 (d, 1H), 2.55 (d, 1H). MS (CI): 251 (M<sup>+</sup>).

Scheme 16 below illustrates the synthesis and structure of 10,11-dihydro- 5-[N-2 methyloxirane]5-H-dibenzo[b,f]azepin (20). 20r, 20s

24



SCHEME 16(20, 20r, 20s)


5 10,11-dihydro-5-[N-β isopropylamino-2propanol)] 5-H-dibenzo  
[b,f]azepin monohydrochloride, 25 bi, 25 bis, 25 bir:

**General procedure:** A suspension of **20** (1.5 g, 5.0 mmol) in *iso*-propanol (50 ml) was warmed to 30 °C, *iso*-propylamine (10 mmol) was added dropwise, the mixture was stirred overnight at this temperature and then allowed to cool to room temperature. The solvent was removed under reduced  
 10 pressure, and the crude product was recrystallized from ethyl acetate at 0 °C.

The precipitation was filtered to give the appropriate free base. The solid was dissolved in toluen and HCl was bubbled untill a precipitation reappeared. The precipitation was filtered and dried under reduced pressure to give 25 bi salt. <sup>1</sup>H NMR (chloroform): 7.07 (bs, 4H), 6.93 (m, 2H), 6.73 (bt,  
 15 2H) 3.79 (m, 1H), 3.15 (s, 4H), 2.78 (m, 1H), 2.72 (d, 2H), 2.55 (d, 2H), 1.01 (d, 6H). MS (CI): 311 (M<sup>+</sup>).

Scheme 17 below illustrates the synthesis and structure of 10,11-dihydro-5-[N-β isopropylamino-2propanol)]5-H-dibenzo[b,f]azepin monohydrochloride, 25 bi, 25 bis and 25 bir.

25



SCHEME 17(25bi, 25bir, 25bis)

Example 4Antiarrhythmic defibrillating activity oftricyclic dibenzazepin and 11-Oxo-dibenzodiazepin compounds

10,11-dihydro-5-[(3-N-alkyl)1-oxopropyl] 5-H-dibenzo[b,f]azepin  
 monohydrochloride, 10,11-dihydro-5-[N-? isopropylamino-2propanol])  
 5-Hdibenzo [b,f]azepin monohydrochloride and 11-Oxo-10,11- dihydro - 5 -  
 10 [(N-alkyl)1 - oxopropyl] - 5 - H-dibenzo [b,e] [1,4] diazepin hydrochloride  
 were synthesized as is described hereinabove and tested for their effectiveness  
 in transforming SVF to TVF, as described herein. The desired outcome was  
 determined to be the ability of the drug molecule to reduce or abolish the  
 occurrence of artificially induced SVF, and its transformation to the  
 15 spontaneously defibrillating TVF. The experimental method was as follows.

Hydrochloride salts of the compounds were dissolved in saline.  
 Activity of each compound was tested in cats of both sexes. Cats were  
 anesthetized with 15-25 mg/kg intravenous sodium pentobarbital. The tested  
 compounds were injected intravenously (1-3 mg/kg). The heart of each  
 20 experimental subject was exposed through midline thoracotomy, and a room  
 air respirator was applied through a tracheal cannula. Lead II  
 electrocardiogram and intra-arterial blood pressure was recorded on a Grass

Polygraph (Grass Instrument Co., Quincy, MA, USA).

Fibrillating stimuli (a train of rectangular pulses of 2 to 15 V, 100 pulses/sec and duration of 0.1 to 1.0 msec, for a period of 1 sec) were delivered through two silver needle electrodes attached to the pericardium on the left ventricle. Fibrillating stimuli were one and a half to twice the strength of the fibrillating threshold.

Animals were designated as having SVF of VF failed to terminate spontaneously within 90 sec, and required electrical defibrillation. Animals that exhibited two to five consecutive episodes of VF of short (20 to 60 sec) duration were designated as having TVF. VF was induced before and after drug administration, according to a previously described procedure [13]. The type of VF was examined before and 2 to 3 min after drug treatment. Each cat served as its own control. In some experiments, a  $\beta$ -adrenergic blocker (propranolol, 0.1-0.6 mg/kg) was administered after drug administration, in order to evaluate the neutralizing effect of the blocker on the compound-induced catecholamine levels. Table 1 below presents the results.

TABLE 1

Compound	Relative activity in dosage of 1 mg/kg	Relative activity in dosage of 2 mg/kg	Relative activity in dosage of 3 mg/kg
15a	50%	67%	100%
15b	50%	75%	100%
15c	64%	67%	83%
15d	44%	67%	100%
15e	0	0	33%
15f	0	0	0
15g	0	0	0
11a	0	-	50%
11b	0	50%	50%
11c	0	0	0
25bi	100%	100%	-
25bir	50%	100%	-
25bis	100%	100%	-

a. TVF /VF (%)

### Example 5

#### Suitable formulations for administration of tricyclic dibenzoazepin and 11-Oxo-dibenzodiazepin compounds

5       The tricyclic dibenzoazepin and 11-Oxo-dibenzodiazepin derivatives of  
the present invention, including free base or salt form, can be administered to a  
subject in a number of ways, which are well known in the art. Hereinafter the  
term "tricyclic dibenzoazepin derivatives" refers to the group of dibenzoazepin  
derivatives in free base form and the group of dibenzoazepin derivatives in a  
10   salt form. Hereinafter the term "tricyclic dibenzodiazepin derivatives" refers to  
the group of dibenzodiazepin derivatives in free base form and the group of  
dibenzodiazepin derivatives in a salt form. Hereinafter, the term "subject"  
refers to the human or any lower animal to which the tricyclic dibenzoazepin  
or 11-Oxo-dibenzodiazepin derivative is administered. For example,  
15   administration may be done topically (including ophtalmically, vaginally,  
rectally, intranasally), orally, or parenterally, for example by intravenous bolus  
or drip, intraperitoneal, subcutaneous, or intramuscular (cardiac) injection.

Formulations for topical administration may include but are not limited  
to lotions, ointments, gels, creams, suppositories, drops, liquids, sprays,  
20   transdermal patches and powders. Conventional pharmaceutical carriers,  
aqueous, powder or oily bases, thickeners and the like may be necessary or  
desirable. In addition to the formulations described previously, a compound of  
the present invention may also be formulated as a depot preparation. Such  
long acting formulations may be administered by implantation (for example  
25   subcutaneously or intramuscularly) or by intramuscular injection. Thus, for  
example, the preparation may be formulated with suitable polymeric or  
hydrophobic materials (for example, as an emulsion in an acceptable oil) or  
ion exchange resins, or as sparingly soluble derivatives such as sparingly  
soluble salts.

Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, sachets, capsules or tablets. Thickeners, diluents, flavorings, dispersing aids, emulsifiers or binders may be desirable.

5 Formulations for parenteral administration may include but are not limited to sterile aqueous solutions, which may also contain buffers, diluents and other suitable additives.

Dosing is dependent on the severity of the symptoms of arrhythmic or fibrillating occurrence and on the responsiveness of the subject to the tricyclic dibenzoazepin derivatives. Persons of ordinary skill in the art can easily  
10 determine optimum dosages, dosing methodologies and repetition rates.

#### Example 6

##### Method of treatment or prevention of ventricular fibrillation

As noted above, the compounds of the present invention, which are  
15 tricyclic dibenzoazepin and 11-Oxo-dibenzodiazepin derivatives, have been shown to be effective defibrillating agents. The following example is an illustration only of a method of treating VF with the dibenzoazepin and 11-Oxo-dibenzodiazepin derivatives, and is not intended to be limiting.

The method includes the step of administering the tricyclic  
20 dibenzoazepin or 11-Oxo-dibenzodiazepin derivatives, in a pharmaceutically acceptable carrier as described in Example 3 above, to a subject to be treated. The tricyclic dibenzoazepin or 11-Oxo-dibenzodiazepin derivative is administered according to an effective dosing methodology, preferably until a predefined endpoint is reached, such as the prevention of VF occurrence or  
25 abnormal cardiac activity. Optionally and preferably, the compound is administered parenterally.

According to another preferred embodiment of the present invention, the compound is used as an adjunct or additive treatment for a patient who has received an implanted defibrillator, such that the compound is administered to  
30 the patient as previously described.

Example 7Method of manufacture of a medicamentcontaining a tricyclic dibenzazepin and 11-Oxo-dibenzodiazepin derivative

5       The following is an example of a method of manufacturing a tricyclic dibenzoazepin and 11-Oxo-dibenzodiazepin derivative. First, the tricyclic dibenzoazepin or 11-Oxo-dibenzodiazepin derivative is synthesized in accordance with good pharmaceutical manufacturing practice. Examples of methods of synthesizing the tricyclic dibenzoazepin and  
10   11-Oxo-dibenzodiazepin derivatives were given previously herein. Next, the tricyclic dibenzoazepin or 11-Oxo-dibenzodiazepin derivative is placed in a suitable pharmaceutical carrier, as described in Example 3 above, again in accordance with good pharmaceutical manufacturing practice.

15       It will be appreciated that the above descriptions are intended only to serve as examples, and that many other embodiments are possible within the spirit and the scope of the present invention.

20       Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.


LIST OF REFERENCES CITED

1. Olshausen K.V., Witt T., Pop T., Treese N., Bethge K., Meyer J. Sudden cardiac death while wearing a Holter monitor. *Am. J. Cardiol.*, 67: 381-386, 1991.
2. Vaughan Williams E.M. Classification of antiarrhythmic drugs. In: Sandoe E., Flasted-Jansen E. Olesen K.H., eds. *Symposium on Cardiac Arrhythmias*. Sudertalje Seden: AB Astra, pp. 449-472, 1970.
3. Sanna G., Archidiacono R. Chemical ventricular defibrillation of the human heart with bretylium tosylate. *Am. J. Cardiol.* 32: 982-986, 1973.
4. CAST Investigators. Preliminary report: Effect of ecainide and flecainide on mortality in a randomized trial of arrhythmia suppression after myocardial infarction. *N. Engl. J. Med.* 321: 406-412, 1989.
5. CAST II Investigators - Effect of the antiarrhythmic agent moricizine on after myocardial infarction. *N. Engl. J. Med.* 327: 227-233, 1992.
6. Manoach M., Varon D., Neuman M., Erez M. Editorial – A new group of defibrillatory drugs in the classification antiarrhythmic agent. *Int. J. Cardiol.* 21: 211-217, 1988.
7. Manoach M., Varon D., Erez M. The role of catecholamines on intracellular coupling, myocardial cell synchronization and self ventricular defibrillation. *Mol. Cell. Biochem.* 147: 181-185, 1995.
8. Manoach M., Erez M., Varon D. Properties required for self ventricular defibrillation: Influence of age and drugs. *Cardiol. Elderly.* 1: 337-344, 1993.
9. Manoach M., Beker B., Erez M., Varon D., Netz H. Spontaneous termination of electrically induced ventricular fibrillation. In: Macfarlane P.W. ed. *Progress in Electrophysiology*. London: Pitman Medical Publications Ltd, pp. 361-365, 1979.
10. Erez M., Varon D. Dibenzazepines and phenothiazines – A winding path to defibrillatory antiarrhythmic agents. *Exp. Clin. Cardiol.* 2: 59-63, 1997.
11. Kaverina N.V., Skoldinov A.P. New cardiovascular drugs among



## WHAT IS CLAIMED IS:


1. A compound having a general formula (II):



wherein  $R_1$  is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and  $C(=O)(CH_2)_nNR'R''$ ,  $(CH_2)_nCHOHCH_2NR'R''$ , wherein  $n$  is an integer;  $R_Q$ ,  $R_T$ ,  $R'$ , and  $R''$  are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and  $t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

2. The compound of claim 1, wherein  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5,  $R'$ ,  $R_Q$ ,  $R_T$ , are each a hydrogen and  $R''$  is as defined above.
3. The compound of claim 2, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of propyl, *n*-butyl, *tert*-butyl and with the proviso that  $R''$  is not a methyl or an ethyl moiety.
4. The compound of claim 2, wherein  $n$  is 1 or 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -cycloalkyl or  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.
5. The compound of claim 4, wherein  $m$  is 1 and  $R''$  is an aromatic 6-member ring.

6. A composition for treating or preventing a cardiac disorder, comprising a pharmaceutically effective amount of a compound in combination with a pharmaceutically acceptable carrier selected from the group consisting of a slow release carrier, an implant and a transdermal patch, said compound being a member of a group having the formula:



wherein,

A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', wherein *n* is an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl, and sulfonylamide; *q* and *t* are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

7. The composition of claim 6, wherein A and B are each a carbon, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are each a hydrogen, and R<sub>1</sub> is C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', *n* being 0-5, R' and R are each hydrogen and R'' is as defined above.

8. The composition of claim 7, wherein *n* is 1 or 2 and R'' is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, and *tert*-butyl.

9. The composition of claim 7, wherein  $n$  is 1 or 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -cycloalkyl or  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.


10. The composition of claim 9, wherein  $m$  is 1 and  $R''$  is an aromatic 6-member ring.

11. The composition of claim 6, wherein  $A$  is  $CR_2R_3$  or  $C=O$  and  $B$  is  $CR_4R_5$ ;  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5,  $R$  and  $R'$  are each hydrogen and  $R''$  is as defined above.

12. The composition of claim 11, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of ethyl, propyl, *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl.

13. The composition of claim 11, wherein  $n$  is 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.

14. A method for treating or preventing a cardiac disorder in a subject, the method comprising the step of administering a pharmaceutically effective amount of a compound, said compound being a member of a group having the formula:



wherein,

$A$  is  $CH$ ,  $CR_2R_3$  or  $C=O$ ;  $B$  is  $CH$ ,  $CR_4R_5$  or  $NR_6$ , wherein  $R_2$ ,  $R_3$ ,  $R_4$ ,

$R_5$  and  $R_6$  are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl; or A and B together are  $C=C$ ;  $R_1$  is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and  $C(=O)(CH_2)_nNR'R''$ ,  $(CH_2)_nCHOHCH_2NR'R''$ , wherein  $n$  is an integer;  $R_Q$ ,  $R_T$ ,  $R'$ , and  $R''$  are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and  $t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

15. The method of claim 14, wherein A and B are each a CH moiety,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5,  $R$  and  $R'$  are each hydrogen and  $R''$  is as defined above.

16. The method of claim 14, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl.

17. The method of claim 15, wherein  $n$  is 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.

18. The method of claim 14, wherein A is  $CR_2R_3$  or  $C=O$  and B is  $CR_4R_5$ ;  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5,  $R$  and  $R'$  are each hydrogen and  $R''$  is as defined above.

19. The method of claim 18, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, and *tert*-butyl


20. The method of claim 18, wherein  $n$  is 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.

21. The method of claim 14, wherein the cardiac disorder is ventricular fibrillation or ischemia.

22. The method of claim 15, wherein said compound is administered to the subject parenterally.

23. The method of claim 15, wherein an implanted defibrillator is implanted in the subject, such that said compound is an adjunct treatment to defibrillation by said implanted defibrillator.

24. A method for stopping the occurrence of ventricular fibrillation in a subject, the method comprising the step of administering a pharmaceutically effective amount of a compound, said compound being a member of a group having the formula:



wherein,

A is CH,  $CR_2R_3$  or  $C=O$ ; B is CH,  $CR_4R_5$  or  $NR_6$ , wherein  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl; or A and B together are  $C=C$ ;  $R_1$  is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and  $C(=O)(CH_2)_nNR'R''$ ,  $(CH_2)_nCHOHCH_2NR'R''$ , wherein  $n$  is an integer;  $R_Q$ ,

$R_T$ ,  $R'$ , and  $R''$  are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and  $t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

25. The method of claim 24, wherein A and B are each a CH moiety,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5,  $R$  and  $R'$  are each hydrogen and  $R''$  is as defined above..

26. The method of claim 25, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, and *tert*-butyl.

27. The method of claim 25, wherein  $n$  is 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.

28. The method of claim 24, wherein A is  $CR_2R_3$  or  $C=O$  and B is  $CR_4R_5$ ;  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are each a hydrogen, and  $R_1$  is  $C(=O)(CH_2)_nNR'R''$ ,  $n$  being 0-5 and  $R'$  is a hydrogen and  $R''$  is as defined above.

29. The method of claim 28, wherein  $n$  is 2 and  $R''$  is an alkyl selected from the group consisting of ethyl, propyl, *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl.


30. The method of claim 28, wherein  $n$  is 2 and  $R''$  is saturated or unsaturated  $(CH_2)_m$ -(hetero)aryl,  $m$  being 0-5.

31. The method of claim 24, wherein said compound is administered

to the subject parenterally.

32. The method of claim 24, wherein an implanted defibrillator is implanted in the subject, such that said compound is an adjunct treatment to defibrillation by said implanted defibrillator.

33. A method of locally treating or preventing a disorder of a tissue of a subject comprising the step of locally applying onto said tissue a composition for treating or preventing a cardiac disorder, said composition comprising a pharmaceutically effective amount of a compound in combination with a pharmaceutically acceptable carrier, said compound being a member of a group having the formula:



wherein A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', wherein *n* is an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl, and sulfonylamide; *q* and *t* are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

34. The method of claim 33, wherein the step of locally applying the

composition onto said tissue further comprises the steps of:

- (i) applying the composition to an implant; and
- (ii) inserting said implant into said tissue.


35. The method of claim 34, wherein said tissue is cardiac tissue.

36. The method of claim 32, wherein the step of locally applying the composition onto said tissue further comprises the steps of:

- (i) applying the composition to a transdermal patch; and
- (ii) applying said patch into said tissue.

37. The method of claim 36, wherein said tissue is skin.

38. A method for treating or preventing a cardiac disorder in a subject, the method comprising the step of inducing cardiac sympathetic activity by administering a compound to the subject, said compound being a member of a group having the formula:



wherein,

A is CH, CR<sub>2</sub>R<sub>3</sub> or C=O; B is CH, CR<sub>4</sub>R<sub>5</sub> or NR<sub>6</sub>, wherein R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted (CH<sub>2</sub>)<sub>m</sub>-(hetero)aryl; or A and B together are C=C; R<sub>1</sub> is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and C(=O)(CH<sub>2</sub>)<sub>n</sub>NR'R'', (CH<sub>2</sub>)<sub>n</sub>CHOHCH<sub>2</sub>NR'R'', wherein *n* is an integer; R<sub>Q</sub>, R<sub>T</sub>, R', and R'' are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or



unsubstituted  $(\text{CH}_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and  $t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof.

39. The method of claim 38, wherein A and B are each a CH moiety,  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $\text{R}_5$  are each a hydrogen, and  $\text{R}_1$  is  $\text{C}(=\text{O})(\text{CH}_2)_n\text{NR}'\text{R}''$ ,  $n$  being 0-5 and R and R' are each hydrogen and R'' is as defined above.

40. The method of claim 39, wherein  $n$  is 2 and R'' is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl.

41. The method of claim 39, wherein  $n$  is 2 and R'' is saturated or unsaturated  $(\text{CH}_2)_m$ -(hetero)aryl,  $m$  being 0-5.

42. The method of claim 38, wherein A is  $\text{CR}_2\text{R}_3$  or  $\text{C}=\text{O}$  and B is  $\text{CR}_4\text{R}_5$ ;  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$ ,  $\text{R}_5$  and  $\text{R}_6$  are each a hydrogen, and  $\text{R}_1$  is  $\text{C}(=\text{O})(\text{CH}_2)_n\text{NR}'\text{R}''$ ,  $n$  being 0-5 and R' is a hydrogen and R'' is as defined above.

43. The method of claim 42, wherein  $n$  is 2 and R'' is an alkyl selected from the group consisting of methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, and *tert*-butyl.

44. The method of claim 42, wherein  $n$  is 2 and R'' is saturated or unsaturated  $(\text{CH}_2)_m$ -(hetero)aryl,  $m$  being 0-5.

45. The method of claim 38, wherein the cardiac disorder is ventricular fibrillation or ischemia.

101069455 (53)

101069455 (53)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION


(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number  
WO 01/15656 A3

- (51) International Patent Classification?: A61K 31/55, C07D 223/18, 243/10, A61P 9/00, 9/10
- (21) International Application Number: PCT/IL00/00510
- (22) International Filing Date: 27 August 2000 (27.08.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 131685 1 September 1999 (01.09.1999) IL
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report
- (88) Date of publication of the international search report: 30 August 2001
- (48) Date of publication of this corrected version: 18 April 2002
- (15) Information about Correction: see PCT Gazette No. 16/2002 of 18 April 2002, Section II
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette*
- (71) Applicant (*for all designated States except US*): TECHNION RESEARCH AND DEVELOPMENT FOUNDATION LTD. [IL/IL]; Senate House, Technion City, 32000 Haifa (IL).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): EREZ, Mordechai [IL/IL]; 19 Shmuel Barkai Street, 69697 Tel Aviv (IL). LEVY, Ofra [IL/IL]; 8 Gedera Street, 49724 Petah Tikva (IL). KEINAN, Ehud [IL/IL]; 8 Moran Street, 23840 Timrat (IL).
- (74) Agent: EHRLICH, Gal; G.E. Ehrlich (1995) Ltd., Bezalel Street 28, 52521 Ramat Gan (IL).

(54) Title: TRICYCLIC COMPOUNDS AND THEIR USES AS ANTIARRHYTHMIC ANTIFIBRILLATORY AND DEFIBRILLATORY AGENTS



(II)

(57) Abstract: A compound having a general formula (II) wherein  $R_1$  is saturated or unsaturated alkyl, amino-alcohol, diamino, cycloalkyl, and  $C(=O)(CH_2)_nNR'R''$ ,  $(CH_2)_nCHOHCH_2NR'R''$ , wherein  $n$  is an integer;  $R_Q$ ,  $R_T$ ,  $R'$ , and  $R''$  are each independently a hydrogen, halogen, hydroxyl, saturated, unsaturated, aliphatic, or branched alkyl, substituted or unsubstituted  $(CH_2)_m$ -(hetero)aryl, and sulfonylamide;  $q$  and

$t$  are each an integer independently selected from 1-4; and pharmaceutically acceptable salts thereof, and the new therapeutic uses thereof and similar compounds as defibrillating, and/or anti-fibrillatory, and/or anti-arrhythmic and/or anti-ischemic drugs.

WO 01/15656 A3

Docket No.  
**02/23383**

# Declaration and Power of Attorney For Patent Application

## English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

### TRICYCLIC COMPOUNDS AND THEIR USES AS ANTIARRHYTHMIC ANTIFIBRILLATORY AND DEFIBRILLATORY AGENTS

the specification of which

☐

is attached hereto.

☒

was filed on 27 August 2000 as ~~United States Application No.~~ or PCT

International Application Number PCT/IL00/00510

and was amended on \_\_\_\_\_

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

131685  
(Number)

IL  
(Country)

01/September/1999  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

I hereby claim the benefit under 35 U.S.C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112. I acknowledge the duty to disclose to the United States Patent and Trademark Office all the information known to me to be material to patentability as defined in Title 37, C.F.R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

PCT/IL00/00510

27 August 2000

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

**SOL SHEINBEIN, Registration Number 25,457**  
**MARTIN MOYNIHAN, Registration Number 40,338**

Send Correspondence to:

**G.E. EHRLICH (1995) LTD.**  
**c/o ANTHONY CASTORINA**  
**2001 JEFFERSON DAVIS HIGHWAY**  
**SUITE 207**  
**ARLINGTON, VIRGINIA 22202, USA**

Direct Telephone Calls to: *(name and telephone number)*

**Anthony Castorina**

**Tel. No. (703) 415-1581**

**Fax No. (703) 415- 4864**

FULL NAME OF SOLE OR FIRST INVENTOR	<b>Mordechai EREZ</b>	
Sole or first inventor's signature	<i>M. Erez</i>	Date <b>2.17.2002</b>
Residence	: 19 Shmuel Barkai Street, 69697 Tel Aviv, Israel	
Citizenship	: Israeli	
Post Office Address	: 19 Shmuel Barkai Street, 69697 Tel Aviv, Israel <b>ILX</b>	

FULL NAME OF SECOND INVENTOR , IF ANY	<b>Ofra LEVY</b>	
Second inventor's signature		Date
Residence	: 8 Gedera Street, 49724 Petach Tikva, Israel	
Citizenship	: Israeli	
Post Office Address	: 8 Gedera Street, 49724 Petach Tikva, Israel	

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

**SOL SHEINBEIN, Registration Number 25,457**  
**MARTIN MOYNIHAN, Registration Number 40,338**

Send Correspondence to: **G.E. EHRLICH (1995) LTD.**  
**c/o ANTHONY CASTORINA**  
**2001 JEFFERSON DAVIS HIGHWAY**  
**SUITE 207**  
**ARLINGTON, VIRGINIA 22202, USA**

Direct Telephone Calls to: *(name and telephone number)*

**Anthony Castorina** **Tel. No. (703) 415-1581**  
**Fax No. (703) 415- 4864**


FULL NAME OF SOLE OR FIRST INVENTOR	<b>Mordechai EREZ</b>
Sole or first inventor's signature	Date _____
Residence	: 19 Shmuel Barkai Street, 69697 Tel Aviv, Israel
Citizenship	: Israeli
Post Office Address	: 19 Shmuel Barkai Street, 69697 Tel Aviv, Israel

FULL NAME OF SECOND INVENTOR, IF ANY	<b>Ofra LEVY</b>
Second inventor's signature	Date <u>18/02/02</u>
Residence	: 8 Gedera Street, 49724 Petach Tikva, Israel
Citizenship	: Israeli
Post Office Address	: 8 Gedera Street, 49724 Petach Tikva, Israel

300

FULL NAME OF THIRD INVENTOR , IF ANY		<b>Ehud KEINAN</b>
Third inventor's signature	<i>Ehud Keinan</i>	Date <u>2-26-02</u>
Residence	: 8 Moran Street, 23840 Timrat, Israel	
Citizenship	: Israeli	
Post Office Address	: 8 Moran Street, 23840 Timrat, Israel <i>DX</i>	

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☒ Page(s) 31 of specification <sup>was</sup> ~~were~~ not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not  
present  
for scanning. (Document title)

☐ *Scanned copy is best available.*